## Indiana State Department of Health MINIMUM IMMUNIZATION REQUIREMENTS FOR SCHOOL ENTRY\* 2010-2011

Grade	Minimum Immunization Requirements
Pre-Kindergarten	<ul> <li>4 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), pediatric diphtheria-tetanus vaccine (DT), or any combination of the three are required.</li> <li>3 doses of either oral polio (OPV) or inactivated polio (IPV) vaccine in any combination.</li> <li>3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).</li> <li>1 dose of measles (rubeola) vaccine on or after the first birthday.</li> <li>1 dose of rubella (German measles) vaccine on or after the first birthday.</li> <li>2 doses of varicella (chickenpox) vaccine on or after the first birthday and separated by 3 months <i>or</i> physician written documentation of history of chickenpox disease, including month and year of disease.</li> </ul>
Kindergarten	<ul> <li>5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).</li> <li>4 doses of any combination of IPV or OPV. The 4<sup>th</sup> dose must be administered on or after the 4<sup>th</sup> birthday, and at least 6 months after the previous dose. (3 doses of all OPV or all IPV are acceptable if the 3<sub>rd</sub> dose was administered on or after the 4<sup>th</sup> birthday, and at least 6 months after the 2<sup>nd</sup> dose).</li> <li>3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be given on or after 24 weeks of age and no earlier than 16 weeks after the 1<sup>st</sup> dose).</li> <li>2 doses of measles (rubeola) vaccine on or after the first birthday.</li> <li>2 doses of rubella (German measles) vaccine on or after the first birthday.</li> <li>1 dose of rubella (chickenpox) vaccine on or after the first birthday and separated by 3 months <i>or</i> physician written documentation of history of chickenpox disease, including month and year of disease.</li> </ul>
Grade 1	<ul> <li>5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).</li> <li>4 doses of any combination of IPV or OPV by age 4-6 (3 doses of all OPV or all IPV are acceptable if the 3<sub>rd</sub> dose was administered on or after the 4th birthday).</li> <li>3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).</li> <li>2 doses of measles (rubeola) vaccine on or after the first birthday.</li> <li>1 dose of rubella (German measles) vaccine on or after the first birthday or physician written documentation of history of chickenpox disease, including month and year of disease.</li> </ul>

## Indiana State Department of Health MINIMUM IMMUNIZATION REQUIREMENTS FOR SCHOOL ENTRY\* 2010-2011

Grade	Minimum Immunization Requirements
Grades 2-5	<ul> <li>5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).</li> <li>4 doses of any combination of IPV or OPV by age 4-6 (3 doses of all OPV or all IPV are acceptable if the 3<sup>rd</sup> dose was administered on or after the 4<sup>th</sup> birthday).</li> <li>3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).</li> <li>2 doses of measles (rubeola) vaccine on or after the first birthday.</li> <li>1 dose of rubella (German measles) vaccine on or after the first birthday or written history of disease. Parental history of chickenpox disease is acceptable proof of immunity. A signed written statement from the parent/guardian indicating month and year of disease is sufficient.</li> </ul>
Grades 6-12	<ul> <li>5 doses of diphtheria-tetanus-acellular pertussis (DTaP), diphtheria-tetanus-pertussis (DTP), or pediatric diphtheria-tetanus vaccine (DT) (4 doses are acceptable if the 4th dose was administered on or after the 4<sup>th</sup> birthday and at least 6 months after the 3<sup>rd</sup> dose).</li> <li>4 doses of any combination of IPV or OPV by age 4-6 (3 doses of all OPV or all IPV are acceptable if the 3<sup>rd</sup> dose was administered on or after the 4<sup>th</sup> birthday).</li> <li>3 doses of Hepatitis B vaccine (3<sup>rd</sup> dose must be on or after 24 weeks of age).</li> <li>2 doses of measles (rubeola) vaccine on or after the first birthday.</li> <li>1 doses of rubella (German measles) vaccine on or after the first birthday.</li> <li>2 doses of varicella (chickenpox) vaccine on or after the first birthday separated by age-appropriate interval or written history of disease. Parental history of chickenpox disease is acceptable proof of immunity. A signed written statement from the parent/guardian indicating month and year of disease is sufficient.</li> <li>1 dose of tetanus-diphtheria-acellular pertussis vaccine (Tdap) given on or after 10 years of age.</li> <li>1 dose of meningococcal conjugate vaccine (MCV4).</li> </ul>

<sup>\*</sup>For children who have delayed immunizations, please refer to the 2010 CDC "Catch-up Immunization Schedule" to determine adequately immunizing doses. All minimum intervals and ages for each vaccination as specified per 2010 CDC guidelines must be met for a dose to be valid. These guidelines can be found at <a href="https://www.cdc.gov/vaccines/recs/schedules/default.htm">www.cdc.gov/vaccines/recs/schedules/default.htm</a>